

Phase transitions and random matrices

G.M.Cicuta

Dept.of Physics, Univ.of Parma,
Viale delle Scienze, 43100 Parma , Italy.
E-mail : cicuta@fis.unipr.it

Abstract

Phase transitions generically occur in random matrix models as the parameters in the joint probability distribution of the random variables are varied. They affect all main features of the theory and the interpretation of statistical models . In this paper a brief review of phase transitions in invariant ensembles is provided, with some comments to the singular values decomposition in complex non-hermitian ensembles.

1 Phase transitions in invariant hermitian ensembles.

Random matrix ensembles have been extensively studied for several decades, since the early works of E.Wigner and F.Dyson, as effective mathematical reference models for the descriptions of statistical properties of the spectra of complex physical systems. In the past twenty years new applications spurred a large literature both in theoretical physics and among mathematicians. Several monographs review different sides of the physics literature of the past few decades, such as [7] , [10], [18] [40] [41], [59] [92] [94] . Their combined bibliography, although very incomplete, exceeds a thousand papers. Sets of lecture notes are [102] [58] [5][34] [85] [66] . The classic reference is Mehta's book [82].

For a long time studies and applications of random matrix theory in large part were limited to the choice of gaussian random variables for the independent entries of the random matrix. This was due both to the dominant role of the normal distribution in probability theory as well as to the nice analytic results which were obtained. Increasingly, in the past two decades, a wide variety of matrix ensembles were considered, where the joint probability distribution for the random entries depends on a number of parameters. As the latter are al-

lowed to change, the generic occurrence of phase transitions emerged.
In this section, I begin by recalling the problem in the easiest case, the invariant ensemble of hermitian matrices .

Let $H = (H_{ij})_{i,j=1,\dots,N}$ be hermitian random matrix with joint probability density for the independent entries

$$\begin{aligned} P(H_{11}, H_{12}, \dots, H_{NN}) dH &= e^{-N \text{Tr } V(H)} dH \bigg/ \int e^{-N \text{Tr } V(H)} dH \quad , \\ V(H) &= \frac{1}{2} a_2 H^2 + \frac{1}{4} a_4 H^4 + \dots + \frac{1}{2p} a_{2p} H^{2p} \quad , \quad a_{2p} > 0 \quad , \\ dH &= \prod_{i < j} (d \text{Re } H_{ij} d \text{Im } H_{ij}) \prod_i dH_{ii} \end{aligned} \quad (1.1)$$

We are interested in the partition function $Z_N(a_2, a_4, \dots, a_{2p})$, the free energy $F_N(a_2, a_4, \dots, a_{2p}) = -\log Z_N(a_2, a_4, \dots, a_{2p})$, the "one point resolvent" $G_N(z)$, the connected correlator $G_N^{(c)}(z_1, z_2)$

$$\begin{aligned} Z_N(a_2, a_4, \dots, a_{2p}) &= \int e^{-N \text{Tr } V(H)} dH \quad , \quad G_N(z) = \frac{1}{N} \text{Tr} < \frac{1}{z - H} > \\ G_N^{(c)}(z_1, z_2) &= < \text{Tr} \frac{1}{z_1 - H} \text{Tr} \frac{1}{z_2 - H} > - < \text{Tr} \frac{1}{z_1 - H} > < \text{Tr} \frac{1}{z_2 - H} > \end{aligned} \quad (1.2)$$

The name invariant ensemble for the ensemble of these hermitian matrices reminds that since the density $P(H_{11}, H_{12}, \dots, H_{NN})$ is invariant under a similarity transformation $H \rightarrow U H U^{-1}$ with arbitrary unitary matrix U , most of the interesting quantities, like those in eqs.(1.2) may be evaluated from the joint probability density of the eigenvalues.

Also important are the monic polynomials $P_n(z) = z^n + 0(z^{n-1})$, orthogonal on the real line with the weight $e^{-N V(z)}$

$$\begin{aligned} \int_{-\infty}^{\infty} P_n(z) P_m(z) e^{-N V(z)} dz &= h_n \delta_{nm} \quad , \\ z P_n(z) &= P_{n+1}(z) + R_n P_{n-1}(z) \quad , \quad R_n = \frac{h_n}{h_{n-1}} > 0 \quad , \\ Z_N(a_2, a_4, \dots, a_{2p}) &= N! (h_0)^N \prod_{n=1}^{N-1} (R_n)^{N-n} \end{aligned} \quad (1.3)$$

One obtains a non-linear recursion relation for the coefficients R_n . For instance if $V(x) = \frac{a_2}{2} x^2 + \frac{a_4}{4} x^4 + \frac{a_6}{6} x^6$ one has the recurrence relation (see [77] , [63] ,

[32])

$$\begin{aligned} \frac{n}{N} = R_n \left[a_2 + a_4 (R_{n-1} + R_n + R_{n+1}) + a_6 (R_{n-1} + R_n + R_{n+1})^2 + \right. \\ \left. + a_6 (R_{n-2}R_{n-1} - R_{n-1}R_{n+1} + R_{n+1}R_{n+2}) \right] \end{aligned} \quad (1.4)$$

occasionally called "pre-string equation" or the Freud equation. One also introduces the set of orthonormal functions $\psi_n(z)$ and the two point kernel $K_N(x, y)$ in terms of which all n -point correlation functions are expressible

$$\psi_n(z) = \frac{1}{\sqrt{h_n}} e^{-NV(z)/2} P_n(z) \quad , \quad K_N(x, y) = \sum_{j=0}^{N-1} \psi_j(x) \psi_j(y) \quad (1.5)$$

If all the coefficients $a_{2k} > 0$ the statistics of the eigenvalues may be evaluated in the limit $N \rightarrow \infty$, see [14], $\lim_{N \rightarrow \infty} G_N(z) = G(z)$ is holomorphic in the complex z plane, except for a segment $(-A, A)$ on the real axis. Furthermore

$$G(z) = \int_{-A}^A d\mu \frac{\rho(\mu)}{z - \mu} \quad , \quad \rho(\mu) = \lim_{N \rightarrow \infty} \frac{1}{N} \text{Tr} < \delta(\mu - H) > = \lim_{N \rightarrow \infty} K_N(\mu, \mu)$$

The limiting density of eigenvalues $\rho(\lambda)$, the one point correlation function, is the unique solution of the integral equation :

$$V'(\lambda) = 2\mathcal{P} \int_{-A}^A d\mu \frac{\rho(\mu)}{\lambda - \mu} \quad , \quad \int_{-A}^A \rho(\lambda) d\lambda = 1 \quad (1.6)$$

$\rho(\lambda) > 0$ on its support $(-A, A)$, it vanishes as a square root at the boundary $\rho(\lambda) \sim (A - |\lambda|)^{1/2}$. Furthermore the coefficients R_n approach a smooth limit $R(\frac{n}{N}) \sim R(x)$ and

$$\begin{aligned} F(a_2, a_4, \dots, a_{2p}) &= N^2 \left[\int_{-A}^A d\lambda \rho(\lambda) V(\lambda) - \int \int_{-A}^A d\lambda d\mu \rho(\lambda) \rho(\mu) \log |\lambda - \mu| + O\left(\frac{1}{N^2}\right) \right] = \\ &= N^2 \left[- \int_0^1 dx (1-x) \log R(x) - \frac{1}{N} \log h_0 + O\left(\frac{1}{N^2}\right) \right] \end{aligned}$$

The free energy $F(a_2, a_4, \dots, a_{2p})$ is analytic in the couplings $(a_2, a_4, \dots, a_{2p})$. The saddle point solution is equivalent to the resummation of the planar graphs, it is equivalent to the solution from the recurrence relations for R_n and to other techniques, such as the loop equations. The orthogonal polynomial technique and the loop equations are superior to evaluate in a systematic way the terms in the series in the parameter $\frac{1}{(N^2)^k}$ corresponding to the resummation of the graphs which are embeddable on orientable surfaces with k handles [101] .

It was also proved that the connected two point correlators exhibit two different forms of universality (that is independence from the set of coefficients $\{a_{2k}\}$ but depend only on the endpoints $\pm A$) :

Global universality. The limiting connected density-density correlator $\rho_c(\lambda, \lambda')$, after smoothing over a scale much larger than the level spacing Δ_N , is [4] [20]

$$\rho_c(\lambda, \lambda') = -\frac{1}{2\pi^2(\lambda - \lambda')^2} \frac{A^2 - \lambda\lambda'}{\sqrt{A^2 - \lambda^2}\sqrt{A^2 - \lambda'^2}} \quad , \quad \lambda \neq \lambda'$$

Local universality. The limiting two point kernel $K(x, y)$ has the sine law [90] for eigenvalues in the bulk of the spectrum, measured in units of Δ_N

$$K_{\text{bulk}}(s, s') = \frac{\sin[\pi(s - s')]}{\pi(s - s')} \quad , \quad s = \lambda/\Delta_N \quad , \quad s' = \lambda'/\Delta_N$$

or the Airy law, close to the tail of the spectrum (the soft edge) [15] [9]

$$K_{\text{soft}}(s, s') = \frac{Ai(s) Ai'(s') - Ai(s') Ai'(s)}{s - s'} \quad , \quad s \sim N^{2/3} \left(\frac{\lambda}{A} - 1 \right)$$

A general derivation of spectral correlators is given in the recent paper by Kanzieper and Freilikher [70]. By following Shohat [97], the recurrence relation for the orthogonal polynomials $P_n(x)$ is turned into an exact second-order differential equation for the orthogonal functions $\psi_n(z)$

$$\frac{d^2\psi_n(\lambda)}{d\lambda^2} - \mathcal{F}_n(\lambda) \frac{d\psi_n(\lambda)}{d\lambda} + \mathcal{G}_n(\lambda) \psi_n(\lambda) = 0$$

Because of the smooth behavior of the coefficients R_n , in the case where the eigenvalues have support on a single segment, the complicated forms $\mathcal{F}_n(\lambda)$, $\mathcal{G}_n(\lambda)$ simplify at large order and the differential equation leads to the global and the local universality results mentioned above.¹

All the above is deeply affected by **phase transitions**.

The limiting eigenvalue density $\rho(\lambda; a_2, a_4, \dots, a_{2p})$ continued to negative values for one or several coefficients a_{2k} (while keeping $a_{2p} > 0$) may not be positive definite. Then the integral equation (1.6) allows new solutions with eigenvalue density positive definite with support on two or more segments of

¹If a logarithmic singularity is present at the origin, the Bessel law is derived

$$K_{\text{origin}}(s, s') = \frac{\pi}{2} (s s')^{1/2} \frac{J_{\alpha+1/2}(\pi s) J_{\alpha-1/2}(\pi s') - J_{\alpha-1/2}(\pi s) J_{\alpha+1/2}(\pi s')}{s - s'}$$

where s and s' are scaled by the level spacing $\Delta_N(0)$ near the spectrum origin, $s = \lambda/\Delta_N(0)$, and $\alpha > -1/2$. For the simpler situations where the orthogonal polynomials are classical the Bessel law had been derived in [22] [87] [53] [88].

the real axis. The correct solution minimizes the free energy. As one explores the space of the parameters, the free energy is evaluated on different saddle point solutions, which may coincide for certain critical values of the parameters. Then the free energy is usually a continuous but not analytic function of the parameters.

The recursion coefficients R_n no longer have a smooth "continuous" limit, which makes more difficult the orthogonal polynomials solution.

The lack of analyticity of the free energy at critical values of the parameters is analogous to phase transitions related to spontaneous symmetry breaking in classical statistical mechanics. For instance, in the simplest case of potential $V(x) = \frac{a_2}{2}x^2 + \frac{a_4}{4}x^4$, first analysed for negative values of a_2 in papers [96] [25], it is convenient to add a linear term, which explicitly breaks the Z_2 symmetry, $V(x) = a_1x + \frac{a_2}{2}x^2 + \frac{a_4}{4}x^4$, next perform the "thermodynamic limit" $N \rightarrow \infty$, finally remove the symmetry breaking term $a_1 \rightarrow 0$, see [62], [26]. This allows the evaluation of the "order parameter" $\langle \text{Tr}H \rangle$

$$\lim_{a_1 \rightarrow 0} \lim_{N \rightarrow \infty} \langle \text{Tr}H \rangle = \text{sign}(a_1) \theta(-a_2 - 2\sqrt{a_4}) f(a_2, a_4)$$

For the simplest case $V(x) = \frac{a_2}{2}x^2 + \frac{a_4}{4}x^4$, it was shown that the correct ansatz for the recursion coefficients R_n , if $a_2 < -2\sqrt{a_4}$ is that the even R_{2n} and odd R_{2n+1} approach two different smooth "continuous" functions [83]. This period-two ansatz requires a little generalization in the case which includes the infinitesimal symmetry breaking term [84] [23] because it leads to recurrence relations

$$z P_n(z) = P_{n+1}(z) + S_n P_n(z) + R_n P_{n-1}(z) \quad , \quad R_n = \frac{h_n}{h_{n-1}} > 0 \quad ,$$

and it explains the origin of the period-two ansatz.

However in the next simplest case, like $V(x) = \frac{a_2}{2}x^2 + \frac{a_4}{4}x^4 + \frac{a_6}{6}x^6$, or higher order polynomials, corresponding to multiple well potentials, which may not be degenerate, the behaviour of the coefficients R_n is erratic and it is difficult to reproduce the results of the saddle point analysis by the orthogonal polynomials [69] [78] [79] [93] [95].

The lines of "phase transition" in the parameter space, related to the continuation to negative values of the coefficient a_{2p} of the monomial of highest order may be found in terms of previously discussed phase transitions by the addition of an infinitesimal monomial ϵx^{2p+2} , see for instance [27] [68].

Connected correlators, when the support of the eigenvalue density is two segments, were shown [6] [1] to have a different form of global universality, involving elliptic integrals. In the case of multicritical behaviour the local universality form has a modified Bessel law [3].

Important recent works seem to be so powerful and comprehensive to solve the above mentioned ambiguities. The Freud equation is expressed in a matrix Lax representation and the semiclassical asymptotics of the functions $\psi_n(z)$ is obtained in the whole complex z plane by solving a matrix Riemann-Hilbert problem, following earlier works [51] [52]. Very useful is the non-linear steepest descent method devised by Deift and Zhou [35] [36]. This set of works [12] [37] [38] [39] not only provide rigorous and more general solutions for methods and ansatzes previously used, but it seem to provide answers also for the models previously left unsolved (like the asymptotics of recurrence coefficients R_n in general cases). The recent and difficult developments will require some time to be exploited by physicists.

Finally, while almost all investigations related to the invariant ensemble of random matrices considered a probability distribution of the form of exponential of a polynomial like in eq.(1.1), with possible addition of logarithmic terms, like the Penner model and the Kontsevich model, there exist probability distributions, still invariant under diagonalization by unitary matrices, which lead to different forms for the connected correlators [2]. Then the classification of universality classes perhaps is not yet complete, even in invariant one-matrix hermitian ensembles.

2 Further matrix ensembles and singular value decomposition in complex non-hermitian random matrices

Random matrix models more general than the hermitian one-matrix invariant ensemble are often more interesting because of the possibility to describe more interesting statistical models. The analytic solution of multi-matrix models both in the "perturbative phase" or in different phases is more complex. While an ensemble of hermitian random matrices describes triangulations of random orientable surfaces, multi-matrix ensembles are suitable to describe models of classical statistical mechanics on a random two-dimensional lattice. After the breakthrough of the Ising model [71] , [13], it was possible to study random walks and loops [43] , $O(N)$ model [55] [75] [76] [45] [46] [44] , the Potts model [72] [33] [105], surfaces with holes [73] , [30], a special case of 8-vertex model [74] [106] , the chiral random matrix which simulate the spontaneously broken phase transition of QCD [103] [89] [98] [100] [65] [11]. Often it was possible to evaluate the critical exponents at phase transition. Multi-matrix models of hermitian matrices are also a good framework for combinatorial problems like the four-color theorem [28] [24] [47] or the enumeration of meanders [42] [81]. Most influential, for quantum field theorists were the phase transitions in models of random unitary matrices, describing one-plaquette of the lattice formulation of

QCD [57] [17] [67] , the saddle-point solution of the one-matrix ensemble [16] , the Witten conjecture of a master field [104].

In this section I shall recall the singular value decomposition, which plays a role in the analysis of models with rectangular random matrices and square complex non-hermitian matrices.

Let $\phi = (\phi_{ij})_{\substack{i=1,\dots,N \\ j=1,\dots,M}}$ be rectangular random matrix with entries ϕ_{ij} real or complex numbers, and joint probability distribution $P(\phi)$ invariant under $\phi \rightarrow U\phi V$, with U unitary of order N , V unitary of order M

$$\begin{aligned}
P(\phi) d\phi &= e^{-N\text{Tr } V(\phi^\dagger \phi)} d\phi \Bigg/ \int e^{-N\text{Tr } V(\phi^\dagger \phi)} d\phi \\
V(\phi^\dagger \phi) &= \frac{1}{2}a_2(\phi^\dagger \phi) + \frac{1}{4}a_4(\phi^\dagger \phi)^2 + \dots + \frac{1}{2p}a_{2p}(\phi^\dagger \phi)^p \quad , \quad a_{2p} > 0 \quad , \\
d\phi &= \prod_{\substack{i=1,\dots,N \\ j=1,\dots,M}} d\phi_{ij} \quad , \quad \text{if } \phi \text{ is real; or} \\
d\phi &= \prod_{\substack{i=1,\dots,N \\ j=1,\dots,M}} (d\text{Re } \phi_{ij} d\text{Im } \phi_{ij}) \quad , \quad \text{if } \phi \text{ is complex}
\end{aligned} \tag{2.1}$$

The hermitian matrices $\phi^\dagger \phi$ and $\phi \phi^\dagger$ are positive semi-definite, have the same non-vanishing eigenvalues $t_i = \sigma_i^2$, where σ_i are the singular values of ϕ , themselves positive definite. It is straightforward to evaluate the statistics of the singular values in the limit $N \rightarrow \infty$, $M \rightarrow \infty$, while the ratio $N/M = L$ is fixed, by the ordinary saddle point analysis. Let us consider first $L \geq 1$. The probability distribution (2.1) is

$$e^{-N \sum_{i=1,\dots,M} V(\sigma_i^2)} J(\sigma_i) \prod_{i=1,\dots,M} d\sigma_i \Bigg/ \int e^{-N \sum_{i=1,\dots,M} V(\sigma_i^2)} J(\sigma_i) \prod_{i=1,\dots,M} d\sigma_i$$

where the Jacobian $J(\sigma_i)$ may be evaluated with help from [60] , [61]

$$\begin{aligned}
J(\sigma_i) &= \prod_{i=1,\dots,M} (\sigma_i)^{N-M} \prod_{1 \leq i < j \leq M} |\sigma_i^2 - \sigma_j^2| \quad , \quad \text{if } \phi \text{ is real,} \\
&= \prod_{i=1,\dots,M} (\sigma_i)^{2N-2M+1} \prod_{1 \leq i < j \leq M} |\sigma_i^2 - \sigma_j^2|^2 \quad , \quad \text{if } \phi \text{ is complex}
\end{aligned}$$

For the simple case $V(\phi^\dagger \phi) = \frac{1}{2}a_2(\phi^\dagger \phi) + \frac{1}{4}a_4(\phi^\dagger \phi)^2$ one easily obtains [29] for complex rectangular matrix ϕ

$$G(z) = \lim_{\substack{N \rightarrow \infty \\ M \rightarrow \infty}} \frac{1}{M} \text{Tr}_M < \frac{1}{z - \phi^\dagger \phi} > = \int_A^B dt \frac{u(t)}{z - t} \quad , \quad 0 \leq A \leq B$$

$$u(t) = \frac{1}{\pi} \sqrt{(B-t)(t-A)} \left(\frac{a_4}{4} + \frac{a_4(A+B)}{8t} + \frac{a_2}{2t} \right) \quad (2.2)$$

The extrema A, B are given by the usual couple of algebraic equation. Then for $a_4 = 0$ one finds the distribution of singular values for rectangular matrices, which is the generalization of Wigner "semi-circle law"

$$u(\sigma) = \frac{a_2}{\pi} \frac{\sqrt{(B-\sigma^2)(\sigma^2-A)}}{\sigma} \quad , \quad \text{for } \sqrt{A} \leq \sigma \leq \sqrt{B}$$

$$A = \left(\frac{\sqrt{L}-1}{a_2} \right)^2 \quad , \quad B = \left(\frac{\sqrt{L}+1}{a_2} \right)^2$$

Returning now to eq.(2.2) for square complex matrices , $L = 1$, one finds a "perturbative" phase for $a_2 > -2\sqrt{a_4}$ with $A = 0$ where observables correspond to resummation of planar graphs, and a "non-perturbative" phase for $a_2 < -2\sqrt{a_4}$ where $A > 0$, quite similar to the random hermitian case. These results were rediscovered by several authors [8] [86] [48] who introduced the hermitian matrix H and the partition function $\mathcal{Z}$

$$H = \begin{pmatrix} 0 & \phi \\ \phi^\dagger & 0 \end{pmatrix} \quad , \quad \mathcal{Z} = \int DH e^{-\beta \text{Tr} V(H^\dagger H)} \sim$$

$$\sim \int_{-\infty}^{\infty} \left[\prod_{i=1}^M dx_i e^{-2\beta V(x_i^2)} \right] \prod_{i=1}^M |x_i|^{2N-2M+1} \prod_{1 \leq i < j \leq M} (x_i^2 - x_j^2)^2$$

The eigenvalues x_i of the "chiral" matrix H are in two to one correspondence with the singular values σ_i of A : $x_i = \pm \sigma_i$. The technique of using the auxiliary matrix H in the study of square complex non-hermitian matrix ϕ was developed by [49] [50] [21] into a powerful method to obtain the distribution $\rho(x, y)$ of complex eigenvalues λ_i , see also the similar and simultaneous paper [64] . Indeed

$$\begin{aligned} \rho(x, y) &= \frac{1}{N} \sum_{i=1}^N \langle \delta(x - \text{Re } \lambda_i) \delta(y - \text{Im } \lambda_i) \rangle = \\ &= \frac{1}{\pi} \frac{\partial}{\partial z} \frac{\partial}{\partial z^*} \frac{1}{N} \langle \text{Tr}_{(N)} \log(z - \phi)(z^* - \phi^\dagger) \rangle = \\ &= \frac{1}{\pi} \frac{\partial}{\partial z} \frac{\partial}{\partial z^*} \frac{1}{N} \left[\langle \text{Tr}_{(2N)} \log H \rangle - i\pi N^2 \right] \end{aligned}$$

where now

$$H = \begin{pmatrix} 0 & \phi - z \\ \phi^\dagger - z^* & 0 \end{pmatrix}$$

Diagrammatic rules may then be used to evaluate the resolvent $\mathcal{G}(\eta; z, z^*)$

$$\mathcal{G}(\eta; z, z^*) = \frac{1}{2N} \langle \text{Tr}_{(2N)} \frac{1}{\eta - H} \rangle = \frac{\eta}{N} \langle \text{Tr}_N \frac{1}{\eta^2 - (z^* - \phi^\dagger)(z - \phi)} \rangle$$

Finally in terms of the integrated density of eigenvalues of H , $\Omega(\mu; z, z^*) = \frac{1}{2N} \langle \text{Tr}_{(2N)} \theta(\mu - H) \rangle$, Feinberg and Zee [50] obtain

$$\rho(x, y) = -\frac{4}{\pi} \int_0^\infty d\mu \frac{\partial}{\partial z} \frac{\partial}{\partial z^*} \frac{\Omega(\mu; z, z^*)}{\mu}$$

As specific examples, Feinberg and Zee consider probability distributions $P(\phi, \phi^\dagger)$ for the complex matrix ϕ of the form (2.1), invariant under $\phi \rightarrow e^{i\alpha} \phi$, $\phi^\dagger \rightarrow e^{-i\alpha} \phi^\dagger$, where it is natural to expect that the distribution of complex eigenvalues $\rho(x, y)$ has rotational symmetry $\rho(x, y) = \rho(r)/(2\pi)$, as it indeed happens with the Ginibre gaussian ensemble [56]. For the simple model with only two coefficients a_2 , a_4 they find that for $a_2 > -2\sqrt{a_4}$ the complex eigenvalues fill (non-uniformly) a disk centered at the origin, while for $a_2 < -2\sqrt{a_4}$ they fill a ring. This is expected from the analogous distribution of the singular values of the matrix ϕ . Next they prove the surprising "single ring theorem" asserting that for a generic polynomial potential of the form (2.1), even in the multiple-well cases where the distribution of the singular values of the matrix ϕ has support on several segments of the positive real axis, the distribution of eigenvalues of the matrix ϕ may only have one ring at most. It seems to me that in such cases the assumption of rotational symmetry should be checked. If it turns out, by using an explicitly symmetry breaking term to be removed after the thermodynamic limit $N \rightarrow \infty$, that one obtains different distributions, dependent on the direction of the symmetry breaking probe, it would be a remarkable example of spontaneous symmetry breaking of rotational symmetry.

3 Conclusion.

Phase transitions generically occur in the study of ensembles of random matrices, as the parameters in the joint probability distribution of the random variables are varied. They are important for the physics interpretation of statistical models and they affect all the main features of random matrix theory. In invariant one-matrix ensembles recent progress of mathematicians seems to solve long standing problems related to multi-cut solutions with no symmetry. Phase transitions in ensembles of complex non-hermitian matrices were recently explored and it is likely that a richer variety of phase transitions will be discovered. Random matrix ensembles with a preferential basis, like band matrices were studied since the beginning of random matrix theory , see for instance [80] [31] [54] [99] . Even the simplest cases of tridiagonal matrices with random site or random hopping could be analytically solved only for a very limited choice of the probability distribution . Well known discrepancies between the moment method and numerical methods suggest the presence of phase transitions which seem more difficult to analyse than in case of invariant ensembles.

References

- [1] G.Akemann, Higher genus correlators for the hermitian matrix model with multiple cuts, Nucl.Phys. **B482** (1996) 403 ; the generalization for singular values of complex matrices is G.Akemann, Universal correlators for multi-arc complex matrix models, Nucl.Phys. **B507** (1997) 475.
- [2] G.Akemann, G.M.Cicuta, L.Molinari, G.Vernizzi, Compact support probability distributions in random matrix theory, Phys.Rev.**E59** (1999) 1489 , and Non-universality of compact support probability distributions in random matrix theory, Phys.Rev.**E60** (1999) 5287-5292.
- [3] G.Akemann, P.H.Damgaard, U.Magnea, S.M.Nishigaki, Multicritical microscopic spectral correlators of hermitian and complex matrices, Nucl.Phys. **B519** (1998) 682.
- [4] J.Ambjorn, J.Jurkiewicz, Y.Makeenko, Multiloop correlators for two-dimensional quantum gravity, Phys.Lett. **B251** (1990) 517 ; the evaluation of correlators in $1/N^2$ expansion was also shown to be universal in J.Ambjorn, L.Chekhov, C.F.Kristjansen, Y.Makeenko, Matrix model calculations beyond the spherical limit, Nucl.Phys. **B404** (1993) 127 and erratum in Nucl.Phys. **B449** (1995) 681 ; and generalized to supermatrices in [91].
- [5] J.Ambjorn, Quantization of geometry, in 1994 Les Houches, ed. F.David, P.Ginsparg, J.Zinn-Justin, North-Holland 1996.
- [6] J.Ambjorn, G.Akemann, New universal spectral correlators, J.Phys. **A29** (1996) L555.
- [7] J.Ambjorn, B.Durhuus, T.Jonsson, Quantum Geometry, Cambridge Univ.Press 1997.
- [8] A.Anderson, R.C.Myers, V.Periwal, Branched polymers from a double-scaling limit of matrix models, Nucl.Phys. **B360** (1991) 463-479.
- [9] E.L.Basor, H.Widom, Determinants of Airy operators and applications to random matrices, J.Stat.Phys. **96** (1999) 1-20.
- [10] C.W.J.Beenakker, Random-matrix theory of quantum transport, Rev.Mod.Phys. **69** (1997) 731.
- [11] Berbenni-Bitsch, M.E.Meyer, S.Schafer, J.J.Verbaarschot, T.Wettig, Microscopic universality in the spectrum of the lattice Dirac operator, Phys.Rev.Lett. **80** (1998) 1146.
- [12] P.Bleher, A.Its, Semiclassical asymptotics of orthogonal polynomials, Riemann-Hilbert problem, and universality in the matrix model, (1997) pag.1-105.

- [13] D.V.Boulatov, V.A.Kazakov, The Ising model on a random planar lattice : the structure of phase transition and the exact critical exponents, Phys.Lett.**B186** (1987) 379.
- [14] A.Boutet de Monvel, L.Pastur, M.Shcherbina, On the statistical mechanics approach in the random matrix theory : integrated density of states, J.Stat.Phys. **79** (1995) 585.
- [15] M.J.Bowick, E.Brezin, Universal scaling of the tail of the density of eigenvalues in random matrix models, Phys.Lett. **B268** (1991) 21.
- [16] E.Brezin, C.Itzykson, G.Parisi, J.B.Zuber , Planar diagrams, Comm. Math.Phys. **59** (1978) 35.
- [17] E.Brezin, D.J.Gross, The external field problem in the large N limit of QCD, Phys.Lett. **B97** (1980) 120.
- [18] E.Brezin, S.R.Wadia, The large N expansion in quantum field theory and statistical physics, World scientific 1993.
- [19] E.Brezin, A.Zee , Universality of the correlations between eigenvalues of large random matrices , Nucl.Phys. **B402** (1993) 613.
- [20] E.Brezin, A.Zee , Correlation functions in disordered systems, Phys.Rev **E 49** (1994) 2588.
- [21] E.Brezin, S.Hikami , A.Zee , Oscillating density of states near zero energy for matrices made of blocks with possible application to the random flux problem, Nucl.Phys. **B464** (1996) 411.
- [22] B.V.Bronk, Exponential ensemble for random matrices, J.Math.Phys. **6** (1965) 228.
- [23] R.C.Brower, N.Deo, S.Jain, C-I Tan, Symmetry breaking in the double-well hermitian matrix models, Nucl.Phys. **B405** (1993) 166-187.
- [24] L.Ckekhov, C.Kristjansen , Hermitian matrix model with plaquette interaction, Nucl.Phys. **B479** (1996) 683.
- [25] G.M.Cicuta, L.Molinari, E.Montaldi, Large N phase transitions in low dimensions, Mod.Phys.Lett.**1** (1986) 125-129.
- [26] G.M.Cicuta, L.Molinari, E.Montaldi, Large- N spontaneous magnetization in zero dimension, J.Phys.**A 20** (1987) L67-L70.
- [27] G.M.Cicuta, L.Molinari, E.Montaldi, Multicritical points in matrix models, J.Phys.**A 23** (1990) L421-L425.

- [28] G.M.Cicuta, L.Molinari, E.Montaldi, Matrix models and graph colouring, Phys.Lett.**B306** (1993) 245.
- [29] G.M.Cicuta, L.Molinari, E.Montaldi, R.Riva , Large rectangular random matrices, J.Math.Phys. **28** (1987) 1716. This work follows the previous investigations A.Barbieri, G.M.Cicuta, E.Montaldi, Nuovo Cimento **A84** (1984) 173 and C.M.Canali, G.M.Cicuta, L.Molinari, E.Montaldi, Nucl.Phys. **B265** (1986) 485.
- [30] G.M.Cicuta, L.Molinari, E.Montaldi, S.Stramaglia, A matrix model for random surfaces with dynamical holes, J. Phys. **A29** (1996) 3769-3785.
- [31] A.Crisanti, G.Paladin, A.Vulpiani, Products of random matrices in statistical physics , Springer series in solid-state sciences 104, Springer-Verlag 1993.
- [32] C.Crnkovic, P.Ginsparg, G.Moore, The Ising model, the Yang-Lee edge singularity and the $2D$ quantum gravity, Phys.Lett.**237B** (1990) 196-201.
- [33] J.M.Daul , Q-states Potts model on a random planar lattice, hep-th/9502014.
- [34] F.David, Simplicial quantum gravity and random lattices, in Gravitation and Quantization, Les Houches 1992 Session LVII, B.Julia and J.Zinn-Justin eds., North Holland 1995.
- [35] P.Deift and X.Zhou, A steepest descent method for oscillatory Riemann-Hilbert problems. Asymptotics for the mKdV equation, Ann. of Math. **137** (1993) 295-370.
- [36] P.Deift and X.Zhou, Asymptotics for the Painleve II equation, Comm.Pure and Appl.Math. **48** (1995) 277-337.
- [37] P.Deift , A.R.Its, X.Zhou, A Riemann-Hilbert approach to asymptotic problems arising in the theory of random matrix models and also in the theory of integrable statistical mechanics, Ann.of Math. **146** (1997) 149-235.
- [38] P.Deift, T.Kriecherbauer, K.T-R McLaughlin, S.Venakides, X.Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, preprint 1-71.
- [39] P.Deift, T.Kriecherbauer, K.T-R McLaughlin, S.Venakides, X.Zhou, Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory, preprint 1-106.

- [40] K.Demeterfi, Two-dimensional quantum gravity, matrix models and string theory, Int.J. Mod.Phys. **A8** (1993) 1185-1244.
- [41] P.Di Francesco, P.Ginsparg, J.Zinn-Justin, 2D gravity and random matrices, Phys.Rep. **254** (1995) 1-133.
- [42] P.Di Francesco, O.Golinelli, E.Guitter, Meander, folding and arch statistics, in Combinatorics and Physics, Mathematical and Computer Modelling 144 (1996).
- [43] B.Duplantier, I.K.Kostov, Geometrical critical phenomena on a random surface of arbitrary genus, Nucl.Phys. **B340** (1990) 491.
- [44] B.Durhuus, C.Kristjansen, Phase structure of the $O(n)$ model on a random lattice for $n > 2$, Nucl.Phys. **B483** (1997) 535-551.
- [45] B.Eynard, C.Kristjansen, Exact solution of the $O(n)$ model on a random lattice, Nucl.Phys. **B455** (1995) 577-618.
- [46] B.Eynard, C.Kristjansen, More on the exact solution of the $O(n)$ model on a random lattice and an investigation of the case $|n| > 2$, Nucl.Phys. **B466** (1996) 463.
- [47] B.Eynard, C.Kristjansen, An iterative solution of the three-colour problem on a random lattice, Nucl.Phys. **B516** (1998) 529.
- [48] J.Feinberg, A.Zee, Renormalizing rectangles and other topics in random matrix theory, cond-mat/9609190.
- [49] J.Feinberg, A.Zee, Non-gaussian non-hermitian random matrix theory: phase transitions and addition formalism, Nucl.Phys. **B501** (1997) 643.
- [50] J.Feinberg, A.Zee, Non-hermitian random matrix theory: method of hermitian reduction, Nucl.Phys. **B504** (1997) 579-608.
- [51] A.S.Fokas, A.R.Its and A.V.Kitaev, The isomonodromy approach to matrix models in 2D quantum gravity, Comm.Math.Phys. **147** (1992) 395-430.
- [52] A.S.Fokas, A.R.Its and A.V.Kitaev, Discrete Painleve equations and their appearance in quantum gravity, Comm.Math.Phys. **142** (1991) 313-344.
- [53] P.J.Forrester, The spectrum of random matrix ensembles, Nucl.Phys. **B402** (1993) 709.
- [54] Y.V.Fyodorov, A.D.Mirlin, Statistical properties of eigenfunctions of random quasi 1D one particle hamiltonians, Int.J.Mod.Phys. **B 8** (1994) 3795-3842.

- [55] M.Gaudin, I.Kostov, $O(n)$ model on a fluctuating planar lattice: some exact results, Phys.Lett.**B220** (1989) 200.
- [56] J.Ginibre, Statistical ensembles of complex, quaternion and real matrices, Jour.Math.Phys. **6** (1965) 440-449.
- [57] D.J.Gross, E.Witten, Possible third-order phase transition in the large- N lattice gauge theory, Phys.Rev. **D21** (1980) 446-453.
- [58] D.J.Gross, T.Piran, S.Weinberg, Two dimensional quantum gravity and random surfaces, World Scientific (1992)
- [59] T.Guhr, A.Muller-Groeling, M.A.Weidenmuller, Random matrix theories in quantum physics : common concepts ,Phys.Rep. **299** (1998) 190.
- [60] S.Helgason , Differential geometry, Lie groups and symmetric spaces, Academic Press (1978).
- [61] L.K.Hua, Harmonic analysis of functions of several complex variables in the classical domains, AMS Providence, Rhode Island (1963).
- [62] E.M.Ilgenfritz, Yu.M.Makeenko, T.V.Shahbazyan, On the relation between many-color QCD and the free Nambu string on the lattice, Phys. Lett. **B172** (1986) 81-85.
- [63] A.R.Its and A.V.Kitaev, Mathematical aspects of the non-perturbative $2D$ quantum gravity, Mod.Phys.Lett.A5 (1990) 2079.
- [64] R.A.Janik, M.A.Nowak, G.Papp, I.Zahed, Nonhermitian random matrix models, Nucl.Phys. **B501** (1997) 603.
- [65] R.A.Janik, M.A.Nowak, G.Papp, I.Zahed, The $U(1)$ problem in chiral random matrix models, Nucl.Phys. **B498** (1997) 313-330.
- [66] R.A.Janik, M.A.Nowak, G.Papp, I.Zahed, Various shades of Blue's functions, 1997 Zakopane lectures, hep-th/9710103.
- [67] J.Jurkiewicz, K.Zalewski, Phase structure of $U(N \rightarrow \infty)$ gauge theory on a two-dimensional lattice for a broad class of variant actions, Nucl.Phys. **B220** (1983) 167-184.
- [68] J.Jurkiewicz, Regularization of one-matrix models, Phys. Lett. **B245** (1990) 178-184.
- [69] J.Jurkiewicz, Chaotic behaviour in one-matrix models, Phys. Lett. **B261** (1991) 260-268.
- [70] E.Kanzieper, V.Freilikher, Spectra of large random matrices: a method of study, (1998) cond-mat/9809365.

- [71] V.A.Kazakov, Ising model on a dynamical planar random lattice: exact solution , Phys.Lett.**A119** (1986) 140-144.
- [72] V.A.Kazakov, Nucl.Phys.(Proc.Suppl.) **B4** (1988) 93.
- [73] V.A.Kazakov, A simple solvable model of quantum field theory of open strings, Phys.Lett.**B237** (1990) 212.
- [74] V.A.Kazakov, P.Zinn-Justin, Two matrix model with *ABAB* interaction, Nucl.Phys. **B546** (1999) 647 .
- [75] I.Kostov, $O(N)$ vector model on a planar random lattice : spectrum of anomalous dimensions , Mod.Phys.Lett.**A4** (1989) 217-226.
- [76] I.Kostov, M.Staudacher , Multicritical phases of the $O(n)$ model on a random lattice, Nucl.Phys. **B384** (1992) 459.
- [77] O.Lichtenfeld, On eigenvalue tunneling in matrix models, Int.J.Mod.Phys. bf **A7** (1992) 2335.
- [78] O.Lichtenfeld, R.Ray, A.Ray, Phase diagram and orthogonal polynomials in multiple well matrix models, Int.J.Mod.Phys. **A6** (1991) 4491-4515.
- [79] O.Lichtenfeld, Semiclassical approach to finite N matrix models, Int.J.Mod.Phys. **A7** (1992) 7097-7118.
- [80] E.H.Lieb, D.C.Mattis, Mathematical physics in one dimension, Academic Press 1966.
- [81] Y.Makeenko, H.Win Pe, Supersymmetric matrix models and the meander problem, (1996) hep-th/9601139.
- [82] M.L.Mehta, Random Matrices, 2nd ed., Academic Press, 1991.
- [83] L.Molinari, Phase structure of matrix models through orthogonal polynomials, J.Phys. **A21** (1988) 1-6.
- [84] L.Molinari, E.Montaldi , The large N magnetization in matrix models revisited, Nuovo Cimento **D15** (1993) 293.
- [85] M.Mulase, Lectures on the asymptotic expansion of a hermitian matrix integral (1998) math-ph/9811023.
- [86] R.C.Myers, V.Periwal, From polymers to quantum gravity: triple scaling in rectangular random matrix models, Nucl.Phys. **B390** (1993) 716-746.
- [87] T.Nagao, K.Slevin, Nonuniversal correlations for random matrix ensembles, J.Math.Phys. **34** (1993) 2075 ; Laguerre ensembles of random matrices : nonuniversal correlation functions, J.Math.Phys. **34** (1993) 2317.

- [88] T.Nagao, P.J.Forrester, Asymptotic correlations at the spectrum edge of random matrices, Nucl.Phys. **B435** (1995) 401.
- [89] M.A.Nowak, J.J.Verbaarschot , I.Zahed, Chiral fermions in the instanton vacuum at finite temperature, Nucl.Phys. **B325** (1989) 581-592.
- [90] L.A.Pastur, M.Shcherbina, Universality of the local eigenvalue statistics for a class of unitary invariant random matrix ensembles, J.Stat.Phys. **86** (1997) 109.
- [91] J.C.Plefka, Iterative solution of the supereigenvalue model, Nucl.Phys. **B444** (1995) 333-352 ; The supereigenvalue model in the double-scaling limit, **B448** (1995) 355-372.
- [92] P.Rossi, M.Campostrini, E.Vicari, The large-N expansion of unitary-matrix models, Phys.Rep. **302** (1998) 143-209.
- [93] M.Sasaki, H.Suzuki, Matrix realization of random surfaces, Phys.Rev **D43** (1991) 4015-4028.
- [94] G.W.Semenoff, R.J.Szabo, Fermionic matrix models, Int.J.Mod.Phys. **A12** (1997) 2135-2292.
- [95] D.Senechal , Chaos in the hermitian one-matrix model, Int.J.Mod.Phys. **A7** (1992) 1491.
- [96] Y.Shimamune, On the phase structure of large N matrix models and gauge models, Phys.Lett.**B108** (1982) 407. The authors of [25] and [26] were unaware of this letter.
- [97] J.Shohat , A differential equation for orthogonal polynomials, Duke Math.J. **5** (1939) 401.
- [98] E.Shuryak, J.J.Verbaarschot, Random matrix theory and spectral sum rules for the Dirac operator in QCD, Nucl.Phys. **A560** (1993) 306-320.
- [99] P.G.Silvestrov, Summing graphs for random band matrices, Phys.Rev. **E55** (1997) 6419-6432.
- [100] Y.A.Simonov, Chiral-symmetry breaking in the disordered QCD vacuum, Phys.Rev. **D43** (1991) 3534-3540.
- [101] The discovery by Gerard 't Hooft of the topological expansion, Nucl.Phys. **B72** (1974) 461 is probably the origin of the interest of quantum field theorists in random matrix theory.
- [102] C.A.Tracy, H.Widom , Introduction to random matrices, in Springer lect. notes in physics 424, ed.G.F.Helminck , Geometric and quantum aspects of integrable systems (1993) 103.

- [103] J.J.Verbaarschot, I.Zahed, Spectral density of the QCD Dirac operator near zero virtuality , Phys.Rev.Lett. **70** (1993) 3852 ; J.J.Verbaarschot, The spectrum of the QCD Dirac operator and chiral random matrix theory : the threefold way, Phys.Rev.Lett. **72** (1994) 2531 ; J.J.Verbaarschot, I.Zahed, Random matrix theory and QCD, Phys.Rev.Lett. **73** (1994) 2288.
- [104] E.Witten, Baryons in the $1/N$ expansion, Nucl.Phys. **B160** (1979) 57-115.
- [105] P.Zinn-Justin, The dilute Potts model on random surfaces, cond-mat/9903385.
- [106] P.Zinn-Justin, The six-vertex model on random lattices, Europhys.Lett. **49** (2000) 15-21.